VOL. 680, NO. 1 JOURNAL OF CHROMATOGRAPHY A SEPTEMBER 30, 1994

CONTENTS

6TH INTERNATIONAL SYMPOSIUM ON HIGH PERFORMANCE CAPILLARY ELECTROPHORESIS, SAN DIEGO, CA, JANUARY 31–FEBRUARY 3, 1994

PART I: VOLUME 680, NO. 1

Foreword
by S. Terabe (Hyogo, Japan) 1

TECHNIQUES AND GENERAL

Terminology and nomenclature in capillary electroseparation systems
by J.H. Knox (Edinburgh, UK) 3

Use of zwitterionic detergents for the separation of closely related peptides by capillary electrophoresis
by K.F. Greve, W. Nashabeh and B.L. Karger (Boston, MA, USA) 15

Wall adsorption of small anions in capillary zone electrophoresis induced by cationic trace constituents of the buffer
by B. Gassner, W. Friedl and E. Kenndler (Vienna, Austria) 25

The use of a microconcentric column in capillary electrophoresis
by C. Fujimoto, H. Matsui, H. Sawada and K. Jinno (Toyohashi, Japan) 33

Capillary zone electrophoresis for the determination of dissociation constants
by S.J. Gluck, (Midland MI, USA) and J.A. Cleveland, Jr. (Indianapolis, IN, USA) 43

Investigation of experimental approaches to the determination of pK_a values by capillary electrophoresis
by S.J. Gluck, (Midland MI, USA) and J.A. Cleveland, Jr. (Indianapolis, IN, USA) 49

Imprinted polymers as antibody mimetics and new affinity gels for selective separations in capillary electrophoresis
by K. Nilsson, J. Lindell, O. Norrlöw and B. Sellergren (Lund, Sweden) 57

Preparation of highly condensed polyacrylamide gel-filled capillaries
by Y. Chen, J.-V. Höltje and U. Schwarz (Tübingen, Germany) 63

Separation of very hydrophobic compounds by hydrophobic interaction electrokinetic chromatography
by E.S. Ahuja and J.P. Foley (Villanova, PA, USA) 73

α,ω-Bis-quaternary ammonium alkanes as effective buffer additives for enhanced capillary electrophoretic separation of glycoproteins
by R.P. Oda, B.J. Madden, T.C. Spelsberg and J.P. Landers (Rochester, MN, USA) 85

Pressurized gradient electro-high-performance liquid chromatography
by B. Behnke and E. Bayer (Tübingen, Germany) 93

Microcolumn sample injection by spontaneous fluid displacement
by H.A. Fishman, R.H. Scheller and R.N. Zare (Stanford, CA, USA) 99

Extended path length post-column flow cell for UV–visible absorbance detection in capillary electrophoresis
by S. Kim, W. Kim and J. Hoon Hahn (Pohang, South Korea) 109

Micellar electrokinetic chromatography using high-molecular surfactants. Use of butyl acrylate–butyl methacrylate–methacrylic acid copolymers sodium salts as pseudo-stationary phases
by H. Ozaki and S. Terabe (Hyogo, Japan) and A. Ichihara (Kyoto, Japan) 117

Novel chiral surfactant for the separation of enantiomers by micellar electrokinetic capillary chromatography
by J.R. Mazzeo, E.R. Grover, M.E. Swartz and J.S. Petersen (Milford, MA, USA) 125

Use of β-cyclodextrin polymer as a chiral selector in capillary electrophoresis
by Z. Aturki and S. Fanali (Rome, Italy) 137

Systematic approach to treatment of enantiomeric separations in capillary electrophoresis and liquid chromatography. I. Initial evaluation using propranolol and dansylated amino acids
by S.G. Penn, G. Liu, E.T. Bergström and D.M. Goodall (York, UK) and J.S. Loran (Kent, UK) 147

Practical aspects of chiral separations of pharmaceuticals by capillary electrophoresis. I. Separation optimization
by A. Guttman and N. Cooke (Fullerton, CA, USA) 157

PHENOLIC COMPOUNDS AND O-HETEROCYCLES

Comparison of micellar electrokinetic chromatography (MEKC) with capillary gas chromatography in the separation of phenols, anilines and polynuclear aromatics. Potential field-screening applications of MEKC
by W.C. Brumley and W.J. Jones (Las Vegas, NV, USA) 163

Optimization of separation selectivity in capillary electrophoresis of flavonoids
by P.G. Pietta, P.L. Mauri, L. Zini and C. Gardana (Milan, Italy) 175

Analysis of isoflavones by capillary electrophoresis
by Z.K. Shihabi, T. Kute, L.L. Garcia and M. Hinsdale (Winston-Salem, NC, USA) 181

CARBOHYDRATES AND GLYCOPROTEINS

Determination of carbohydrates in fruit juices by capillary electrophoresis and high-performance liquid chromatography
by A. Klockow (Basle and Zurich, Switzerland), A. Paulus and V. Figueiredo (Basle, Switzerland), R. Amadò (Zurich, Switzerland) and H.M. Widmer (Basle, Switzerland) 187

Application of capillary ion electrophoresis and ion chromatography for the determination of O-acetate groups in bacterial polysaccharides
by R.W. Hepler and C.C. Yu Ip (West Point, PA, USA) 201

High-performance capillary electrophoresis of O-glycosidically linked sialic acid-containing oligosaccharides in glycoproteins as their alditol derivatives with low-wavelength UV monitoring
by K. Kakehi, A. Susami, A. Taga, S. Suzuki and S. Honda (Higashi-osaka, Japan) 209

Analysis of the microheterogeneity of the glycoprotein chorionic gonadotropin with high-performance capillary electrophoresis
by D.E. Morbeck, B.J. Madden and D.J. McCormick (Rochester, MN, USA) 217

Determination of the number and distribution of oligosaccharide linkage positions in O-linked glycoproteins by capillary electrophoresis
by P.L. Weber (Sioux City, 1A and Lawrence, KS, USA) and C.J. Bramich and S.M. Lunte (Lawrence, KS, USA) 225

Separation and analysis of cyclodextrins by capillary electrophoresis with dynamic fluorescence labelling and detection
by S.G. Penn (York, UK) and R.W. Chiu and C.A. Monnig (Riverside, CA, USA) 233

ORGANIC ACIDS

Capillary zone electrophoretic determination of organic acids in cerebrospinal fluid from patients with central nervous system diseases
by A. Hiraoka and J. Akai (Tokyo, Japan), I. Tominaga and M. Hattori (Chiba, Japan) and H. Sasaki and T. Arato (Tokyo, Japan) 243

Determination of organic acids in urine by capillary zone electrophoresis
by M. Shirao, R. Furuta, S. Suzuki and H. Nakazawa (Tokyo, Japan), S. Fujita, (Gokohama-shi, Japan) and T. Maruyama (Osaka, Japan) 247

AMINES, AMINO ACIDS AND PEPTIDES

Enantiomeric resolution of primary amines by capillary electrophoresis and high-performance liquid chromatography using chiral crown ethers
by Y. Walbroehl and J. Wagner (Strasbourg, France) 253

Dual electrochemical detection of cysteine and cystine in capillary zone electrophoresis
by B.L. Lin, L.A. Colón and R.N. Zare (Stanford, CA, USA) 263

Simultaneous detection of thiols and disulfides by capillary electrophoresis–electrochemical detection using a mixed-valence ruthenium cyanide-modified microelectrode
by J. Zhou, T.J. O'Shea and S.M. Lunte (Lawrence, KS, USA) 271

Improvement of laser-induced fluorescence detection of amino acids in capillary zone electrophoresis
by J. Mattusch and K. Dittrich (Leipzig, Germany) 279

Capillary electrophoretic determination of amino acids with indirect absorbance detection
by Y.-H. Lee and T.-I. Lin (Taipei, Taiwan) 287

Analysis of dansyl amino acids in feedstuffs and skin by micellar electrokinetic capillary chromatography
by S. Michaelsen (Tjele, Denmark) and P. Møller and H. Sørensen (Frederiksberg, Denmark) 299

Separation of 24 dansylamino acids by capillary electrophoresis with a non-ionic surfactant
by N. Matsubara and S. Terabe (Hyogo, Japan) 311

Optical resolution of amino acid derivatives by micellar electrokinetic chromatography with N-dodecanoyl-L-serine
by K. Otsuka, K. Karuhaka and M. Higashimori (Osaka, Japan) and S. Terabe (Hyogo, Japan) 317

Simulation and optimization of peptide separation by capillary electrophoresis
by A. Cifuentes and H. Poppe (Amsterdam, Netherlands) 321

Multiple-buffer-additive strategies for enhanced capillary electrophoretic separation of peptides
by R.P. Oda, B.J. Madden, J.C. Morris, T.C. Spelsberg and J.P. Landers (Rochester, MN, USA) 341

PART II: VOLUME 680, NO. 2

PROTEINS

Capillary electrophoresis–matrix-assisted laser-desorption ionization mass spectrometry of proteins
by W. Weinmann (Konstanz, Germany and Research Triangle Park, NC, USA), C.E. Parker, L.J. Deterding and D.I. Papac (Research Triangle Park, NC, USA), J. Hoyes (Manchester, UK), M. Przybylski (Konstanz, Germany) and K.B. Tomer (Research Triangle Park, NC, USA) 353

Evaluation of a novel hydrophilic derivatized capillary for protein analysis by capillary electrophoresis–electrospray mass spectrometry
by R.B. Cole and J. Varghese (New Orleans, LA, USA) and R.M. McCormick and D. Kadlecek (Sunnyvale, LA, USA) 363

Ultra-fast high-performance capillary sodium dodecyl sulfate gel electrophoresis of proteins
by K. Benedek (Thousand Oaks, CA, USA) and A. Guttman (Linz, CA, USA) 375

Control of the cultivation process of antithrombin III and its characterization by capillary electrophoresis
by O.-W. Reif and R. Freitag (Hannover, Germany) 383

Affinity capillary electrophoresis applied to the studies of interactions of a member of heat shock protein family with an immunosuppressant
by J. Liu and K.J. Volk (Wallingford, CT, USA), M.S. Lee (Princeton, NJ, USA), E.H. Kerns (Wallingford, CT, USA) and I.E. Rosenberg (Princeton, Wallingford, CT, USA) 395

Determination of antigen–antibody affinity by immuno-capillary electrophoresis
by N.H.H. Heegaard (Copenhagen, Denmark) 405

Rapid characterization of soy protein and hydrolysates by capillary electrophoresis
by T.M. Wong, C.M. Carey and S.H.C. Lin (St. Louis, MO, USA) 413

Characterization of charge-modified and fluorescein-labeled antibody by capillary electrophoresis using laser-induced fluorescence. Application to immunoassay of low level immunoglobulin A
by F.-T.A. Chen (Fullerton, CA, USA) 419

Characterization of digoxigenin-labeled B-phycoerythrin by capillary electrophoresis with laser-induced fluorescence. Application to homogeneous digoxin immunoassay
by F.-T.A. Chen and S.L. Pentoney, Jr. (Fullerton, CA, USA) 425

Determination of isoelectric points of acidic and basic proteins by capillary electrophoresis
by Y.J. Yao, K.S. Khoo, M.C.M. Chung and S.F.Y. Li (Singapore, Singapore) 431

High-performance capillary electrophoresis for characterization of hapten–protein conjugates used for production of antibodies against soyasaponin I
by H. Frøkiær (Lyngby, Denmark), P. Møller and H. Sørensen (Frederiksberg, Denmark) and S. Sørensen (Lyngby, Denmark) 437

Capillary electrophoresis of the scrapie prion protein from sheep brain
by M.J. Schmerr, K.R. Goodwin and R.C. Cutlip (Ames, IA, USA) 447

Monitoring enzymic peroxidation of NAD^+ by capillary electrophoresis
by V. Dolník and L. Skurský (Brno, Czech Republic) 455

NUCLEIC ACIDS CONSTITUENTS AND PCR PRODUCTS

Analysis of nucleotide pools in human lymphoma cells by capillary electrophoresis
by X. Shao, K. O'Neill, Z. Zhao, S. Anderson, A. Malik and M. Lee (Provo, UT, USA) 463

Quantitative capillary gel electrophoresis assay of phosphorothioate oligonucleotides in pharmaceutical formulations
by G.S. Srivatsa, M. Batt, J. Schuette, R.H. Carlson, J. Fitchett, C. Lee and D.L. Cole (Carlsbad, CA, USA) 469

Capillary gel electrophoresis of oligonucleotides: prediction of migration times using base-specific migration coefficients
by Y. Cordier and O. Roch (Strasbourg, France), P. Cordier (Romansviller, France) and R. Bischoff (Strasbourg, France) 479

Detection of DNA fragments separated by capillary electrophoresis based on their native fluorescence inside a sheath flow
by D.A. McGregor and E.S. Yeung (Ames, IA, USA) 491

High-speed and high-accuracy DNA sequencing by capillary gel electrophoresis in a simple, low cost instrument. Two-color peak-height encoded sequencing at 40°C
by H. Lu, E. Arriaga, D.Y. Chen and N.J. Dovichi (Edmonton, Canada) 497

Activation energy of single-stranded DNA moving through cross-linked polyacrylamide gels at 300 V/cm. Effect of temperature on sequencing rate in high-electric-field capillary gel electrophoresis
by H. Lu, E. Arriaga, D.Y. Chen, D. Figeys and N.J. Dovichi (Edmonton, Canada) 503

Effects of sample matrix and injection plug on dsDNA migration in capillary gel electrophoresis
by M.J. van der Schans, J.K. Allen, B.J. Wanders and A. Guttman (Fullerton, CA, USA) 511

Analysis of polymerase chain reaction-product by capillary electrophoresis with laser-induced fluorescence detection and its application to the diagnosis of medium-chain acyl-coenzyme A dehydrogenase deficiency
by H. Arakawa, K. Uetanaka, M. Maeda and A. Tsuji (Tokyo, Japan) and Y. Matsubara and K. Narisawa (Sendai, Japan) 517

Analysis of DNA restriction fragments and polymerase chain reaction products by capillary electrophoresis
by P.E. Williams, M.A. Marino, S.A. Del Rio, L.A. Turni and J.M. Devaney (Gaithersburg, MD, USA) 525

DYES AND HETEROCYCLIC COMPOUNDS

Determination of synthetic food dyes by capillary electrophoresis
by S. Suzuki, M. Shirao, M. Aizawa and H. Nakazawa (Tokyo, Japan) and K. Sasa and H. Sasagawa (Osaka, Japan) 541

Behaviour of substituted aminomethylphenol dyes in capillary isoelectric focusing with electroosmotic zone displacement
by J. Caslavska and S. Molteni (Berne, Switzerland), J. Chmelík, K. Šlais and F. Matulík (Brno, Czech Republic) and W. Thormann (Berne, Switzerland) 549

Determination of heterocyclic compounds by micellar electrokinetic capillary chromatography
by C. Bjergegaard, H. Simonsen and H. Sørensen (Frederiksberg, Denmark) 561

Separation of charged and neutral isotopic molecules by micellar electrokinetic chromatography in coated capillaries
by M. Chiari, M. Nesi, G. Ottolina and P.G. Righetti (Milan, Italy) 571

DRUGS

Determination of pharmaceuticals and related impurities by capillary electrophoresis
by C.L. Ng, C.P. Ong, H.K. Lee and S.F.Y. Li (Singapore, Singapore) 579

Analysis of structural specificity in antibody–antigen reactions by capillary electrophoresis with laser-induced fluorescence detection
by R.A. Evangelista and F.-T.A. Chen (Fullerton, CA, USA) 587

Determination of components in propolis by capillary electrophoresis and photodiode array detection
by H. Chi, A.K. Hsieh, C.L. Ng, H.K. Lee and S.F.Y. Li (Singapore, Singapore) 593

Capillary electrophoretic chiral separations using cyclodextrin additives. III. Peak resolution surfaces for ibuprofen and homatropine as a function of the pH and the concentration of β-cyclodextrin
by Y.Y. Rawjee, R.L. Williams and G. Vigh (College Station, TX, USA) 599

Optimization of a capillary electrophoresis method to determine the chiral purity of a drug
by E.C. Rickard and R.J. Bopp (Indianapolis, IN, USA) 609

Mixed polymer networks in the direct analysis of pharmaceuticals in urine by capillary electrophoresis
by H. Soini (Bloomington, IN, USA), M.-L. Riekkola (Helsinki, Finland) and M.V. Novotny (Bloomington, IN, USA) 623

ENVIRONMENTAL ANALYSIS AND IONS

Capillary liquid chromatography–mass spectrometry and micellar electrokinetic chromatography as complementary techniques in environmental analysis
by W.C. Brumley, C.M. Brownrigg and A.H. Grange (Las Vegas, NV, USA) 635

Environmental monitoring of wastewater using capillary ion electrophoresis
by S.A. Oehrle (Milford, MA, USA) and R.D. Blanchard, C.L. Stumpf and D.L. Wulfeck (Highland Heights, KY, USA) 645

Simultaneous detection and quantitation of sodium, potassium, calcium and magnesium in ocular lenses by high-performance capillary electrophoresis with indirect photometric detection
by H. Shi, R. Zhang, G. Chandrasekher and Y. Ma (Kirksville, MO, USA) 653

Separation and indirect detection by capillary zone electrophoresis of ppb (w/w) levels of aluminum ions in solutions of multiple cations
by W.R. Barger, R.L. Mowery and J.R. Wyatt (Washington, DC, USA) 659

Capillary electrophoretic separation of organophosphonic acids using borate esterification and direct UV detection
by W.H. Robins and B.W. Wright (Richland, WA, USA) 667

Determination of anions with capillary electrophoresis and indirect ultraviolet detection
by M.M. Rhemrev-Boom (Emmen, Netherlands) 675

AUTHOR INDEX 685